

AD-A112 254


BRANDEIS UNIV WALTHAM MA DEPT OF CHEMISTRY F/8 7/4
STRUCTURE AND SOLID-STATE REACTIVITY OF A NEW POLYMORPH OF TETR--ETC(U)
DEC 81 B M FOXMAN, H MAZUREK N00014-76-C-0822

UNCLASSIFIED

NL

1 1
A 1 1
A 1 1

END
DATE
FILMED
4-82
DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

DTIC FILE COPY

ADA112254

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 8	2. GOVT ACCESSION NO. (12) NR-A112 254	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Structure and Solid-State Reactivity of a New Polymorph of Tetrakis(4-vinylpyridine)-diisothiocyanatocobalt(II)		5. TYPE OF REPORT & PERIOD COVERED Technical Report
7. AUTHOR(s) Bruce M. Foxman and Harry Mazurek		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Brandeis University Department of Chemistry Waltham, Massachusetts 02254		8. CONTRACT OR GRANT NUMBER(s) N00014-76-C-0822
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, VA 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 053-619
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE 1 December 1981
		13. NUMBER OF PAGES 22
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES To be published in Inorganica Chimica Acta		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Polymorphism Coordination compounds Solid-state reaction Phase transitions Crystal structure		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The crystal structure of $\text{Co}(\text{NCS})_2[4\text{-vinylpyridine}]_4$ has been determined by X-ray diffraction methods. The complex crystallizes in space group Pbca, with $a=32.293(15)$, $b=11.416(6)$, $c=17.033(10)$ Å; $z=8$. Full matrix least-squares refinement of positional and thermal parameters for all atoms with 1458 reflections led to $R=0.075$ and $R_w=0.079$. The <u>trans</u> -octahedral complex consists of four vinylpyridine ligands, arranged in a propeller fashion, and two isothiocyanate ligands. Powder diffraction of the thermal decomposition		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 68 IS OBSOLETE
S/N 0102-014-6601

Unclassified

82 03 22 107
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

product, $\text{Co}(\text{NCS})_2(4\text{-vinylpyridine})_2$ indicates an isometric cell, with $a=13.77\text{\AA}$. The analogous $\text{Ni}(\text{II})$ complexes are isomorphous with the $\text{Co}(\text{II})$ complexes, suggesting that the yellow-green complex $\text{Ni}(\text{NCS})_2(4\text{-vinylpyridine})_2$, previously thought to be polymeric, is in fact tetrahedral.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution	
Availability Codes	
Dist	



Unclassified

OFFICE OF NAVAL RESEARCH

Contract No. N00014-76-C-0822

Task No. NR 053-619

TECHNICAL REPORT No. 8

The Structure and Solid-State Reactivity of a New
Polymorph of Tetrakis(4-vinylpyridine)diisothiocyanatocobalt (II)

by

Bruce M. Foxman and Harry Mazurek

Prepared for Publication

in

Inorganica Chimica Acta

Brandeis University

Department of Chemistry

Waltham, Massachusetts 02254

December 1, 1981

Reproduction in whole or in part is permitted for
any purpose of the United States Government
This document has been approved for public release
and sale; its distribution is unlimited.

THE STRUCTURE AND SOLID-STATE REACTIVITY OF
A NEW POLYMORPH OF
TETRAKIS(4-VINYLPYRIDINE)DIISOTHIOCYANATOCOBALT(II)

by

Bruce M. Foxman* and Harry Mazurek

Contribution from the Department of Chemistry
Brandeis University, Waltham, Massachusetts 02254

* Author to whom correspondence should be addressed

ABSTRACT

The crystal structure of $\text{Co}(\text{NCS})_2[4\text{-vinylpyridine}]_4$ has been determined by X-ray diffraction methods. The complex crystallizes in space group Pbca , with $a = 32.293(15) \text{ \AA}$, $b = 11.416(6) \text{ \AA}$, $c = 17.033(10) \text{ \AA}$, and $Z = 8$. Full-matrix least-squares refinement of positional and thermal parameters for all atoms (H atoms fixed at calculated positions), with 1458 reflections ($F > 3.92 \sigma(F)$) led to $R = 0.075$ and $R_w = 0.079$. The trans-octahedral complex consists of four vinylpyridine ligands, arranged in a propeller-shape, and two isothiocyanate ligands. Powder diffraction of $\text{Co}(\text{NCS})_2(4\text{-vinylpyridine})_2$ indicates an isometric cell, with $a = 13.77 \text{ \AA}$. The analogous $\text{Ni}(\text{II})$ complexes are isomorphous with the $\text{Co}(\text{II})$ complexes, suggesting that the yellow-green complex $\text{Ni}(\text{NCS})_2(4\text{-vpy})_2$, previously thought to be polymeric, is in fact tetrahedral.

Introduction

In organic solid-state polymerization reactions, the constraints imposed by the crystal lattice may promote the formation of highly oriented, crystalline product [1]. Single crystals of an organic monomer contain a basic repeat unit, e.g., a carbon-carbon multiple bond, which is an inherent part of the reactant. In transition metal solid-state chemistry, the use of complexes with polyfunctional ligands may allow a close analogy to be drawn to the organic species. For example, we have shown that square-planar nickel(II) complexes of tris (2-cyanoethyl)phosphine polymerize in the solid state to produce polymer crystals with "single crystal" diffraction properties [2], and also undergo polymorphic transitions [3]. We have endeavored to extend our study of such processes to metal complexes with other polyfunctional ligands.

It has been demonstrated that cobalt(II) and nickel(II) complexes of 4-vinylpyridine undergo a series of interesting solid-state reactions [4,5], and also have the ability to act as preferential clathrates [6]. The reactivity of these complexes, i.e. $M(\text{substituted pyridine})_4X_2 \xrightarrow[\Delta]{-2py} M(\text{substituted pyridine})_2X_2$, where X = halogen or pseudohalogen group, is initially similar to that of the analogous pyridine, lutidine, picoline and quinoline complexes [7]. Dichlorobis(4-vinylpyridine)cobalt(II) exists in two forms: an α , low density, blue form [8] and a β , high density,

violet form [9]. The α form is tetrahedrally coordinated while the β form is octahedrally coordinated via bridging chlorine atoms. Upon heating, the vinyl groups of the α form polymerize. However, the vinyl groups in the β form do not polymerize; rather, the β form undergoes a solid-state reaction to produce the α form.

Extensive differential scanning calorimetry and optical microscopic studies have been carried out on the solid-state reaction of $\text{Co(4-vinylpyridine)}_4(\text{NCS})_2$ [5]. In an attempt to comment on the molecular mechanism of the transformation $\text{Co(4-vpy)}_4(\text{NCS})_2 \xrightarrow{\Delta} \text{Co(4-vpy)}_2 + 2(4\text{-vpy})$ we synthesized a previously uncharacterized polymorph of $\text{Co(4-vpy)}_4(\text{NCS})_2$. The present work is a study of the preparation and molecular structure of that material and its decomposition reaction in the solid state.

Experimental

Stoichiometric amounts of Co(NCS)_2 (Alfa/Ventron) and 4-vinylpyridine (Aldrich) are mixed in absolute ethanol; red crystals form upon standing for ten hours. $\text{Co(NCS)}_2(4\text{-vinylpyridine})_2$ can be obtained by prolonged heating (~ 6 hr.) of the red crystals at 115°C , upon which they become blue. The analogous nickel complexes (which are isomorphous) can be formed similarly; a color change from blue to yellow-green is noted.

Collection and Reduction of Diffraction Data. Preliminary Weissenberg ($h0l, h1l$) photographs showed the symmetry and characteristic absences $0k\bar{l}$, $k=2n+1$; $h0\bar{l}$, $l=2n+1$; $hk0$, $h=2n+1$ of the

space group Pbca. A crystal was mounted on a Supper No. 455 goniometer and optically centered on a Syntex P2₁ diffractometer. Most operations were carried out as described previously [10]; other operations are described below. Details of the structure analysis, in outline form, are presented in Table I.

Solution and Refinement. The analysis was performed on a Syntex XTL structure determination system [11]. The analytical scattering factors of Cromer and Waber were used [12a]; real and imaginary components of anomalous scattering were included in the calculations for all nonhydrogen atoms [12b]. The cobalt atom position was found from a three-dimensional Patterson map. Subsequent difference Fourier syntheses and structure factor calculations revealed the remaining atomic positional parameters.

On the final cycles of least-squares refinement, only the Co and S were assigned anisotropic temperature factors; H atoms were fixed at calculated positions ($r_{C-H} = 0.95 \text{ \AA}$) with temperature factors one unit higher than that of the carbon atom to which they were bonded. At convergence, [$\Delta/\sigma_{\max} \leq 0.31$], a weighting scheme analysis revealed no systematic dependence of $w[|F_o| - |F_c|]^2$ on $|F_o|$, $(\sin \theta/\lambda)$, parity of indices, or sequence number. Table II lists the positional and isotropic temperature factors for all atoms.

Supplementary Data Available. Tables of anisotropic thermal parameters, H atom coordinates, and observed vs. calculated structure factors are available on request from the Editor or the authors.

Results and Discussion

The molecular structure of the octahedral complex is depicted in Figure 1. The Co-N(isothiocyanate) bond lengths, 2.100(13) Å and 2.049(12) Å, are comparable to the 2.086(7) Å bond length found in the low-density form of $\text{Co}(\text{NCS})_2(4\text{-vinylpyridine})_4$, reportedly an "unsolvated" or desorbed clathrate complex [13]. The Co-N-C-(isothiocyanate linkage) angles, however, 173.0(11)° and 166.4(12)°, are nearer to linearity than those in the desorbed form (161.4(7)°). The geometrical arrangement of the vinylpyridine rings with respect to one another and to the plane defined by the four N(pyridine) atoms is comparable to that found in the desorbed form, nickel 4-methylpyridine complexes [14], and other tetrakis (pyridine or substituted pyridine) complexes [15]. The vinylpyridine rings are arranged in a staggered "propeller" configuration. The rings trans to one another are nearly orthogonal (N(3) ring 108° to N(4) ring; N(5) ring 94° to N(6) ring). There are no intermolecular contacts less than the sum of the corresponding van der Waals radii, either in the present crystal structure or in that of the low-density polymorph [13]. However, as might be expected, the intermolecular contacts are shorter and more numerous in the high-density polymorph. Table III lists other pertinent bond lengths and angles.

Thermal decomposition of $\text{Co}(4\text{-vpy})_4(\text{NCS})_2$. -- Agnew and Brown [5] have reported that one can generate $\text{Co}(4\text{-vpy})_2(\text{NCS})_2$ by the liberation of 4-vinylpyridine from the analogous tetrakis-(4-vinylpyridine) complex. The former complex is blue, and is similar

to other tetrahedral $\text{Co}(\text{substituted pyridine})_2\text{X}_2$ complexes. We note that the decomposition produces polycrystalline $\text{Co}(4\text{-vpy})_2(\text{NCS})_2$, and all diffraction lines from the powder pattern of $\text{Co}(4\text{-vpy})_2(\text{NCS})_2$ can be indexed to an isometric system [16] with $a = 13.77 \text{ \AA}$ (see Table IV). The density of this presumably tetrahedral complex, $1.46(1) \text{ g cm}^{-3}$ (measured by neutral buoyancy in chloroform-hexane), indicates six molecules per unit cell. This high-symmetry crystal structure need not be disordered; for example, $\text{Co}(4\text{-vpy})_2(\text{NCS})_2$, with $Z = 6$ in space group $P23$, could occupy any of the sites 6f through 6i [17]. Studies in this laboratory of the thermal decomposition of $\text{Ni}(4\text{-vpy})_4(\text{NCS})_2$ (isomorphous with $\text{Co}(4\text{-vpy})_4(\text{NCS})_2$) indicate that the $\text{Ni}(4\text{-vpy})_2(\text{NCS})_2$ product is also cubic, with $a = 13.77 \text{ \AA}$. Thus, the polymeric octahedral structure proposed for $\text{Ni}(4\text{-vpy})_2(\text{NCS})_2$ [4] based on the C-N stretching frequency, and magnetic susceptibility measurements, is inconsistent with our X-ray data.

For $\text{Ni}(4\text{-vpy})_2(\text{NCS})_2$, the infrared spectrum of the yellow-green material produced by decomposition of $\text{Ni}(4\text{-vpy})_4(\text{NCS})_2$ is nearly identical to the published spectrum for the green solution-prepared product [4]. In view of this, and considering the similar thermal behavior as well as the isomorphism between analogous Ni and Co complexes, it appears likely that $\text{Ni}(\text{vpy})_2(\text{NCS})_2$ is tetrahedral.

Polymorphism and Reactivity of $\text{Co}(4\text{-vpy})_4(\text{NCS})_2$. As mentioned above, there are at least two polymorphic forms of the title complex.

We have studied the orthorhombic form, space group Pbca, with $a = 32.293$, $b = 11.416$ and $c = 17.033$ Å; the tetragonal form [13], space group $I4_1/a$, has $a = 17.05$ and $c = 25.79$ Å. While the near-equality of the 17 Å axes in the two polymorphs might suggest a spatial relationship, we have been unable to discover it. This is consistent with the large disparity in crystal densities ($\rho_{\text{orth}} = 1.26$, $\rho_{\text{tetr}} = 1.05$ g-cm⁻³). These structural differences suggested to us that a comparative study of the thermal loss of two vinylpyridine ligands would be worthwhile. Unfortunately, we were unable to prepare the low-density form of the complex by the literature syntheses or alternative methods. All efforts produced only the orthorhombic form of the crystals.

Acknowledgement. -- This work was supported in part by the Office of Naval Research.

References

1. G. Wegner, Pure Appl. Chem., 49, 443 (1977); H. Nakanishi, W. Jones, J. M. Thomas, M. Hasegawa and W. L. Rees, Proc. R. Soc. Lond., A369, 307 (1980); R. H. Baughman, J. Polym. Sci., Polym. Phys. Ed., 12, 1511 (1974).
2. K. Cheng and B. M. Foxman, J. Am. Chem. Soc., 99, 8102 (1977); K. Cheng, B. M. Foxman and S. W. Gersten, Mol. Cryst. Liq. Cryst., 52, 77 (1979).
3. B. M. Foxman, P. L. Goldberg and H. Mazurek, Inorg. Chem., 20, xxxx (1981).
4. N. H. Agnew, R. J. Collin and L. F. Larkworthy, J. Chem. Soc. Dalton, 272 (1974) and references therein.
5. N. H. Agnew and M. E. Brown, J. Polym. Sci.: Polym. Chem. Ed., 12, 1493 (1974) and references therein.
6. W. D. Schaeffer, W. S. Dorsey, D. A. Skinner and C. G. Christian, J. Am. Chem. Soc., 79, 5870 (1957); M. I. Hart and N. O. Smith, J. Am. Chem. Soc., 84, 1816 (1962).
7. L. C. Lewis and N. Fogel, J. Chem. Soc. A, 1141 (1970); E. Jona, V. Jesenak, T. Sramko and J. Gazo, J. Thermal Anal., 2, 389 (1973); D. H. Brown and R. T. Richardson, J. Inorg. Nucl. Chem., 35, 755 (1973).
8. L. J. Admiraal and G. Gafner, Chem. Commun., 1221 (1968).
9. M. Laing and E. Horsfield, Chem. Commun., 902 (1969).
10. B. M. Foxman, Inorg. Chem., 17, 1932 (1978); B. M. Foxman and H. Mazurek, Inorg. Chem., 18, 113 (1979).

11. "Syntex XTL Operations Manual," 2nd ed.; Syntex Analytical Instruments, Cupertino, Calif., 1976.
12. "International Tables for X-ray Crystallography," Kynoch Press, Birmingham, England, 1974; Vol IV, (a) pp 99-101; (b) pp 148-150.
13. G. D. Andreetti and P. Sgarabotto, Cryst. Struct. Commun., 1, 55 (1972).
14. I. S. Kerr and D. J. Williams, Acta Cryst., B33, 3589 (1977); G. D. Andreetti, G. Bocelli and P. Sgarabotto, Cryst. Struct. Commun., 1, 51 (1972).
15. G. J. Long and P. J. Clarke, Inorg. Chem. 17, 1394 (1978); D. J. Hamm, J. Bordner and A. F. Schreiner, Inorg. Chim. Acta, 7, 637 (1973); I. Sotofte and S. E. Rasmussen, Acta Chem. Scand. 21, 2028 (1967).
16. L. V. Azaroff and M. J. Buerger, "The Powder Method in X-ray Crystallography," McGraw-Hill, New York, 1958, pp 61-62.
17. "International Tables for X-ray Crystallography," Vol. I, Kynoch Press, Birmingham, England, 1969.

(D) Refinement,^d with 1458 Data for which $F > 3.92 \sigma(F)$

Weighting of reflections: $w = [\sigma^2(|F_o|) + (p|F_o|)^2]^{-1}$; $p = 0.030$

Isotropic refinement, all nonhydrogen atoms: $R = 0.106$; $R_w = 0.115$

Anisotropic refinement of Co, S(1), S(2), isotropic refinement of the remaining nonhydrogen atoms, fixed contributions from the hydrogen atoms: $R = 0.075$; $R_w = 0.079$

Structure factor calcn, all 3259 data: $R = 0.190$; $R_w = 0.092$

Standard deviation of an obsv of unit weight (SDU): 1.5120

Final difference Fourier map: random peaks $\leq 0.48 \text{ e/\AA}^3$

^aMeasured by flotation in $\text{C}_2\text{H}_5\text{OH}-\text{CHCl}_3$.

$$^b R_s = \sum \sigma(|F_o|) / \sum |F_o|.$$

^cChurchill, M. R.; Lashewycz, R. A.; Rotella, F. J. Inorg Chem., **16**, 265 (1977).

$$^d R = \sum (|F_o| - |F_c|) / \sum |F_o|; R_w = \{ \sum w[|F_o| - |F_c|]^2 / \sum w|F_o|^2 \}^{1/2}.$$

SDU = $\{ \sum w[|F_o| - |F_c|]^2 / (m-n) \}^{1/2}$ where $m(=1458)$ is the number of observations and $n(=172)$ is the number of parameters.

Table II

Atomic Coordinates and Isotropic Temperature Factors^a

ATOM	X	Y	Z	U _{iso}
CO	0.133879(63)	0.07132(15)	-0.05968(11)	0.0571(12)
S(1)	0.19088(17)	0.20270(36)	-0.30521(27)	0.0978(37)
S(2)	0.09133(19)	-0.11650(46)	0.19030(34)	0.1350(49)
N(1)	0.15557(37)	0.13940(95)	-0.16592(74)	0.0694(37)
N(2)	0.10900(34)	0.0029(10)	0.04190(69)	0.0653(36)
N(3)	0.16006(39)	-0.09956(96)	-0.08951(67)	0.0702(39)
N(4)	0.10583(36)	0.2406(10)	-0.03372(74)	0.0690(37)
N(5)	0.07777(40)	0.01386(97)	-0.12270(73)	0.0735(38)
N(6)	0.19946(33)	0.11585(86)	0.00009(64)	0.0528(32)
C(1)	0.17051(43)	0.1667(11)	-0.22593(88)	0.0569(43)
C(2)	0.10230(46)	-0.0497(13)	0.09709(94)	0.0714(48)
C(3)	0.14087(47)	-0.1999(13)	-0.07128(87)	0.0778(49)
C(4)	0.15378(55)	-0.3096(14)	-0.10178(98)	0.0886(55)
C(5)	0.18592(51)	-0.3135(13)	-0.15382(94)	0.0762(50)
C(6)	0.20669(48)	-0.2133(13)	-0.16972(86)	0.0774(50)
C(7)	0.19364(49)	-0.1089(12)	-0.13628(88)	0.0684(46)
C(8)	0.20330(75)	-0.4276(22)	-0.1966(14)	0.1482(97)
C(9)	0.1876(10)	-0.5086(27)	-0.1908(17)	0.223(14)
C(10)	0.08916(54)	0.3070(15)	-0.0893(10)	0.0973(61)
C(11)	0.07136(55)	0.4191(16)	-0.0739(11)	0.1073(59)
C(12)	0.07292(52)	0.4628(13)	-0.0035(11)	0.0852(54)
C(13)	0.09123(55)	0.3980(14)	0.0515(11)	0.0946(56)
C(14)	0.10765(46)	0.2872(12)	0.03597(93)	0.0696(48)
C(15)	0.05687(73)	0.5986(20)	0.0239(14)	0.1452(95)
C(16)	0.03742(91)	0.6451(24)	-0.0186(17)	0.194(12)
C(17)	0.03927(55)	0.0179(13)	-0.09119(97)	0.0801(50)
C(18)	0.00662(54)	-0.0470(14)	-0.1190(11)	0.0928(56)
C(19)	0.01015(55)	-0.1156(13)	-0.19535(97)	0.0829(53)
C(20)	0.04784(54)	-0.1163(12)	-0.22022(86)	0.0725(48)
C(21)	0.08006(48)	-0.0519(13)	-0.19041(91)	0.0754(48)
C(22)	-0.03083(78)	-0.1867(21)	-0.2089(14)	0.1494(88)
C(23)	-0.02311(92)	-0.2665(26)	-0.2371(19)	0.217(14)
C(24)	0.21178(46)	0.2102(12)	-0.01604(93)	0.0630(43)
C(25)	0.24899(47)	0.2373(12)	0.01530(93)	0.0626(41)
C(26)	0.26733(45)	0.1648(12)	0.06738(87)	0.0660(44)
C(27)	0.24453(46)	0.0632(12)	0.09552(78)	0.0691(43)
C(28)	0.20645(42)	0.0433(10)	0.05195(82)	0.0567(40)
C(29)	0.30733(57)	0.1895(14)	0.1049(10)	0.0895(56)
C(30)	0.32971(60)	0.1240(16)	0.1445(12)	0.1179(70)

^a Standard deviations in the least significant digit appear in parentheses in this and subsequent tables. For atoms refined anisotropically, $U_{iso} = U_{eq} = 1/3 \sum_i \sum_j U_{ij} a_i^* a_j^* a_i \cdot a_j$.


Table III
Selected Bond Lengths (\AA) and Angles (deg)

Co-N(1)	2.100(13) \AA	Co-N(1)-C(1)	173.0(11)°
N(1)-C(1)	1.17(2)	N(1)-C(1)-S(1)	179.2(13)
C(1)-S(1)	1.56(2)	Co-N(2)-C(2)	166.4(12)
Co-N(2)	2.049(12)	N(2)-C(2)-S(2)	175.6(14)
N(2)-C(2)	1.14(2)	N(1)-Co-N(2)	177.8(5)
C(2)-S(2)	1.65(2)	N(3)-Co-N(4)	178.3(4)
Co-N(3)	2.197(11)	N(5)-Co-N(6)	175.8(4)
Co-N(4)	2.169(12)	C(5)-C(8)-C(9)	120.1(26)
Co-N(5)	2.185(13)	C(12)-C(15)-C(16)	119.6(23)
Co-N(6)	2.147(11)	C(19)-C(22)-C(23)	110.8(25)
C(5)-C(8)	1.60(3)	C(26)-C(29)-C(30)	129.3(17)
C(8)-C(9)	1.06(4)		
C(12)-C(15)	1.60(3)		
C(15)-C(16)	1.16(4)		
C(19)-C(22)	1.60(3)		
C(22)-C(23)	1.06(4)		
C(26)-C(29)	1.47(2)		
C(29)-C(30)	1.24(3)		

Table IV
Powder Pattern of $\text{Co}(\text{NCS})_2(4\text{-vinylpyridine})_2$

Number	Intensity	hkl
1	95	220
2	50	222
3	60	400
4	70	422
5	50	432, 520
6	100	600, 442
7	40	730
8	80	650, 643
9	20	820, 644
10	20	664
11	20	864, 1004
12	20	880
13	20	884, 1200
14	20	888

Figure I. Molecular structure of the complex.



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 1211 Research Triangle Park, N.C. 27709	1
ONR Western Regional Office Attn: Dr. R. J. Marcus 1030 East Green Street Pasadena, California 91106	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Eastern Regional Office Attn: Dr. L. H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
The Assistant Secretary of the Navy (RE&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
Mr. A. M. Anzalone Administrative Librarian PLASTE/ARRADCOM Bldg 3401 Dover, New Jersey 07801	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
		Mr. James Kelley DTNSRDC Code 2803 Annapolis, Maryland 21402	1

TECHNICAL REPORT DISTRIBUTION LIST, 053

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. R. N. Grimes Department of Chemistry University of Virginia Charlottesville, Virginia 22901	1	Dr. B. Foxman Department of Chemistry Brandeis University Waltham, Massachusetts 02154	1
Dr. M. F. Hawthorne Department of Chemistry University of California Los Angeles, California 90024	1	Dr. T. Marks Department of Chemistry Northwestern University Evanston, Illinois 60201	1
Dr. D. B. Brown Department of Chemistry University of Vermont Burlington, Vermont 05401	1	Dr. J. Zuckerman Department of Chemistry University of Oklahoma Norman, Oklahoma 73019	1
Dr. D. Venezky Chemistry Division Naval Research Laboratory Code 6130 Washington, D.C. 20375	1	Professor O. T. Beachley Department of Chemistry State University of New York Buffalo, New York 14214	1
Dr. J. Adcock Department of Chemistry University of Tennessee Knoxville, Tennessee 37916	1	Professor K. M. Nicholas Department of Chemistry Boston College Chestnut Hill, Massachusetts 02167	1
Dr. A. Cowley Department of Chemistry University of Texas Austin, Texas 78712	1	Professor R. Neilson Department of Chemistry Texas Christian University Fort Worth, Texas 76129	1
Dr. W. Hatfield Department of Chemistry University of North Carolina Chapel Hill, North Carolina 27514	1	Professor M. Newcomb Texas A&M University Department of Chemistry College Station, Texas 77843	1
Dr. D. Seyferth Department of Chemistry Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1	Professor Richard Eisenberg Department of Chemistry University of Rochester Rochester, New York 14627	1
Dr. M. H. Chisholm Department of Chemistry Indiana University Bloomington, Indiana 47401	1		

